

KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY

NEOPRINIL® POUR-ON

For the treatment and control of internal and external parasites of beef and dairy cattle (including lactating cows) and internal parasites of deer.

ACTIVE CONSTITUENTS

5 g/L eprinomectin.

INDICATIONS

For the treatment and control in all ages of beef and dairy cattle (including lactating cows) of:

- **Adult and immature gastrointestinal worms:** *Ostertagia ostertagi* (including inhibited immature), *Ostertagia lyrata* (adult only), *Haemonchus contortus*, *Trichostrongylus axei*, *Trichostrongylus colubriformis*, *Oesophagostomum radiatum*, *Nematodirus helvetianus*, *Bunostomum phlebotomum*, *Trichuris discolor* (adult only), *Trichuris globulosa* (adult only), *Cooperia oncophora*, *Cooperia punctata* and *Cooperia surnabada*.
- **Lungworm:** Adult and immature *Dictyocaulus viviparus*.
- **Sucking and biting lice:** *Bovicola bovis*, *Linognathus vituli*, *Haematopinus euryternus* and *Solenopotes capillatus*.
- **Sarcoptic and chorioptic mites:** *Sarcoptes scabiei* var *bovis* and *Chorioptes bovis*.

And for the treatment and control in all ages of deer of:

- **Adult and immature gastrointestinal worms:** *Ostertagia circumcincta*, *Ostertagia leptospicularis*, *Trichostrongylus axei* and *Oesophagostomum venulosum*.
- **Lungworms:** *Dictyocaulus ekerti* (*viviparus*).

PERSISTENT ACTIVITY IN CATTLE

NEOPRINIL® POUR-ON for cattle and deer when given at the recommended dosage of 1 ml/10 kg liveweight prevents reinfection of cattle with the following internal parasites for the time period indicated.

- *Ostertagia* spp.: 28 days
- *Cooperia* spp.: 21 days
- *Nematodirus helvetianus*: 14 days
- *Oesophagostomum radiatum*: 28 days
- *Dictyocaulus viviparus*: 28 days
- *Haemonchus contortus*: 14 days
- *Trichostrongylus* spp.: 14 days

For best results NEOPRINIL® POUR-ON for cattle and deer should be part of a total parasite control program. Check the effectiveness of anthelmintics at least annually. Follow recommended parasite management practices to reduce the potential for resistance to develop. Consult your veterinarian for a complete parasite control program.

DOSAGE AND ADMINISTRATION

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

The dose is 1ml for each 10 kg of liveweight to give 0.5 mg eprinomectin per kg liveweight. For pour-on application only. Apply along the topline in a narrow strip extending from the withers to the tailhead. Do not apply to areas of the backline covered with mud or manure. See pack for dose rate guide. A representative sample of animals should be weighed before treatment either with scales or a weighband. Dose rate to be based on heaviest cattle in each group (bulls, cows, steers, calves etc). Do not underdose. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing. Correct drenching technique should be used. Heavier animals (for example mature bulls) should receive an additional 5 ml for each 50 kg over 650 kg. Rainfall before or after treatment will not affect efficacy.

Ask your **Virbac Area Sales Manager** or **veterinarian** for more information

WITHHOLDING PERIODS

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

- Meat (cattle): Nil
- Bobby calves: Nil
- Meat (deer): 7 days
- Milk (cattle): Nil

PRECAUTIONS

Keep this and all drugs out of reach of children. If accidental skin contact occurs, wash the affected area immediately with soap and water. If accidental eye exposure occurs, flush immediately with water and get medical attention if irritation persists.

WARNING

Hazard classification: **6.8B, 9.1A, 9.2C, 9.3C, 9.4A.** Harmful - may cause developmental damage to the unborn child from repeated oral exposure. Keep out of reach of children. Do not handle until all safety precautions have been read and understood. Ecotoxic - very toxic to aquatic organisms. Avoid contamination of any water supply with product or empty container. Harmful to the soil environment. Harmful to terrestrial vertebrates. Very toxic to terrestrial invertebrates. Apply only as directed.



DISPOSAL

Preferably dispose of product by use. Otherwise dispose of product and packaging at an approved landfill/facility. Triple rinse empty container. Spread rinsate onto waste ground or pasture. Recycle empty rinsed container at an approved recycling centre. Otherwise puncture or crush empty container and dispose of at an approved landfill/ facility. Do not use container for any other purpose.

EMERGENCY MANAGEMENT

In case of spillage wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Sweep up solids without creating dust. Wash area with water and absorb with further inert material. Dispose of waste safely. Do not carry more than 1 litre on a passenger service vehicle.

STORAGE

Close container when not in use. Store at or below 25°C. Protect from light. Stores containing equal to or more than 100 litres of NEOPRINIL® POUR-ON require bunding and are subject to signage. See the Hazardous Substances Emergency Management Regulations 35-41, Identification Regulations 34 & 35 and NZS 8409 for further information.

Registered pursuant to the ACVM Act 1997, No. A11278. See www.foodsafety.govt.nz for registration conditions.

Registered to and distributed in New Zealand by:

VIRBAC NEW ZEALAND LIMITED
26-30 Maui Street, Pukete,
Hamilton 3200, New Zealand.
Phone: 0800 VIRBAC (0800 847 222)
Visit us at nz.virbac.com

Shaping the future
of animal health

